

Properties of functional differential equation in Banach spaces

Diana Otrocol and Veronica Ana Ilea

T. Popoviciu Institute of Numerical Analysis, Romanian Academy, Cluj-Napoca, Romania,
Babeş-Bolyai University, Department of Mathematics, Cluj-Napoca, Romania
email: dotrocol@ictp.acad.ro, vdarzu@math.ubbcluj.ro

In this paper we consider a functional differential equations with abstract Volterra operator. The equations involving abstract Volterra operators have been investigated by many authors ([?], [?], [?]) and the references therein).

The aim of our paper is to extend the results from [?] to Banach spaces and to consider some applications of this equations in bounded domain, complex space and L_2 space.

References

- [1] C. Corduneanu, *Abstract Volterra equations (a survey)*, Math. and Comp. Modelling, **32** (2000), 1503-1528.
- [2] D. Otrocol, *Abstract Volterra operators*, Carpathian J. Math., **24** (2008) no. 3, 370-377.
- [3] D. Otrocol, V.A. Ilea, *Qualitative properties of functional differential equation*, Electr. J. Qual. Th. Diff. Eq., 2014, No. 47, 1–8.
- [4] M.A. Şerban, I.A. Rus, and A. Petruşel, *A class of abstract Volterra equations, via weakly Picard operators technique*, Math. Inequal. Appl., **13** (2010), 255-269.